

Kinematics of Extragalactic Radio Sources

A Model for Symmetric Jets

M. Thulasidas¹

New Initiatives, Institute for Infocomm Research, 21 Heng Mui Keng Terrace, Singapore 119613.
e-mail: manoj@i2r.a-star.edu.sg

Submitted to A&A: November 29, 2004; AA/2004/2445

Abstract. Symmetric extragalactic radio sources exhibit a baffling array of features that are very poorly understood. What is the origin of the remarkable symmetry between the two lobes? Why is the emission in the radio frequency (RF) range? Why are the jets so well collimated, implying long term memory of the core? Why do the core regions emit blue and ultraviolet light? What is the origin of transient X-ray and γ ray bursts? In this article, we present a model for the kinematics of extragalactic radio sources that explains these features. First, we show that the traditional explanation (Rees 1966) for the apparent superluminal motion (which is based on the light travel time effect) requires the sources to be highly asymmetrical. This is in stark contrast with the observation of most radio sources with double lobed structures, which tend to look very symmetrical. We establish that an apparent superluminal motion in *both* the jets of these radio sources (galactic or extragalactic) necessarily implies *real* superluminal motion. The light travel time effect influences the way we perceive superluminal motion. An object, moving across our field of vision at superluminal speeds, will appear to us as two objects receding from a single point. Based on this effect, we derive the kinematical properties of extragalactic radio sources and explain the puzzling features listed above. Furthermore, we derive the time evolution of the hot spots and compare it to the proper motions reported in the literature (Biretta et al. 1999). We also compare the time evolution of a microquasar (Mirabel & Rodríguez 1999) with our model and show excellent agreement. This model can also explain the observed blue/UV spectrum (and its time evolution) of the core region and the RF spectrum of the lobes, and why the radio sources appear to be associated with galactic nuclei. We make other quantitative predictions, which can be verified.

Key words. radio continuum: galaxies – galaxies: active – galaxies: jets – galaxies: kinematics and dynamics – X-rays: bursts – Gamma rays: bursts

1. Introduction

Transverse velocities of a celestial object can be measured almost directly using angular measurements. The angular rate can be translated to a speed using the known (or estimated) distance of the object from us. In the past few decades, scientists have observed (Biretta et al. 1999; Zensus 1997) objects moving at apparent transverse velocities significantly higher than the speed of light. Fig. 1 shows one such example. It shows the image of the galaxy M87 from the Space Telescope Science Institute. The top panel is an image from the Hubble space telescope showing a 5000 light year long jet from the galaxy's nucleus. The panel below shows a sequence of Hubble images showing highly superluminal motion, with the slanting lines tracking the moving features (Biretta et al. 1999). Some such superluminal objects were detected within our own galaxy (Mirabel & Rodríguez 1994, 1999; Gisler 1994; Fender et al. 1999).

The special theory of relativity (Einstein 1905) states that nothing can accelerate past the speed of light. A direct measurement of superluminal objects emanating from a single point (or a small region) would be a violation of the special theory of relativity at a fundamental level. However, before proclaiming a contradiction based on the measurement of an apparent superluminal motion, one has to establish that the measurement is not an artifact of the way one perceives transverse velocities. Rees (1966) offered an explanation why such apparent superluminal motion is not in disagreement with the special theory of relativity, even before the phenomenon was discovered. When an object travels at a high speed towards an observer, at a shallow angle with respect to his line of sight, it can appear to possess superluminal speeds. Thus, a measurement of superluminal transverse velocity by itself is not an evidence against the special theory of relativity. In this article, we re-examine this explanation (also known also as the light travel time effect.) We will show that the apparent symmetry of the extragalactic radio sources is inconsistent with this explanation.

Send offprint requests to: M. Thulasidas




Fig. 1. Sequence of Hubble images showing apparent motion at six times the speed of light in the galaxy M87. Photo Credit: John Biretta, Space Telescope Science Institute.




Fig. 2. False colour image of the radio jet and lobes in the hyperluminous radio galaxy Cygnus A. Red shows regions with the brightest radio emission, while blue shows regions of fainter emission. Image courtesy of NRAO/AUI.

Transverse superluminal motions are usually observed in quasars and microquasars. Different classes of such objects associated with Active Galactic Nuclei (AGN) were found in the last fifty years. Fig. 2 shows the radio galaxy, Cygnus A (Perley et al. 1984) – one of the brightest radio objects. Many of its features are common to most extragalactic radio sources – the symmetric double lobes, an indication of a core, an appearance of jets feeding the lobes and the hot spots. Owsianik & Conway (1998) and Polatidis et al. (2002) have reported more detailed kinematical features, such as the proper motion of the hot spots in the lobes. We will show that our per-

ception of an object crossing our field of vision at a constant superluminal speed is remarkably similar to a pair of symmetric hot spots departing from a fixed point with a decelerating rate of angular separation. We will make other quantitative predictions that can be verified, either from the current data or with dedicated experiments.

2. Traditional Explanation




Fig. 3. Illustration of the traditional explanation for the apparent superluminal motion. An object expanding at a speed $\beta = 0.8$, starting from a single point S. The solid circle represents the boundary one second later. The observer is far away on the right hand side, O ($x \rightarrow \infty$). The dashed ellipse is the apparent boundary of the object, as seen by the observer.

First, we look at the traditional explanation of the apparent superluminal motion by the light travel time effect. Fig. 3 illustrates the explanation of apparent superluminal motion as described in the seminal paper by Rees (1966). In this figure, the object at S is expanding radially at a constant speed of $0.8c$, a highly relativistic speed. The part of the object expanding along the direction V_1 , close to the line of sight of the observer, will appear to be travelling much faster. This will result in apparent transverse velocity that can be superluminal.

The apparent speed β' of the object depends on the real speed β and the angle between its direction of motion and the observer's line of sight, θ . As shown in the appendix on Mathematical Details,

$$\beta' = \frac{\beta}{1 - \beta \cos \theta} \quad (1)$$

Fig. 3 is a representation of equation (1) as $\cos \theta$ is varied over its range. It is the locus of β' for a constant $\beta = 0.8$, plotted against the angle θ . The predicted shape of the apparent speed is in complete agreement with what was predicted in 1966 (Fig. 1 in that article (Rees 1966)).

For a narrow range of θ , the transverse component of the apparent velocity ($\beta' \sin \theta$) can appear superluminal. From equation (1), it is easy to find this range:

$$\frac{1 - \sqrt{2\beta^2 - 1}}{2\beta} < \cos \theta < \frac{1 + \sqrt{2\beta^2 - 1}}{2\beta} \quad (2)$$

Thus, for appropriate values of $\beta (> \frac{1}{\sqrt{2}})$ and θ (as given in equation (2)), the transverse velocity of an object can seem su-

perluminal, even though the real speed is in conformity with the special theory of relativity.

While equations (1) and (2) explain the apparent transverse superluminal motion the difficulty arises in the recessional side. Along directions such as V_2 in Fig. 3, the apparent velocity is always smaller than the real velocity. The jets are believed to be emanating from the same AGN in opposite directions. Thus, if one jet is in the $\cos \theta$ range required for the apparent superluminal motion (similar to V_1), then the other jet has to be in a direction similar to V_2 . Along this direction, the apparent speed is necessarily smaller than the real speed, due to the same light travel time effect that explains the apparent superluminal motion along V_1 . Thus, the observed symmetry of these objects is inconsistent with the explanation based on the light travel time effect. Specifically, superluminality can never be observed in both the jets (which, indeed, has not been reported so far). However, there is significant evidence of near symmetric outflows (Laing et al. 1999) from a large number of objects similar to Fig. 2.

One way out of this difficulty is to consider hypothetical superluminal speeds for the objects making up the apparent jets. Note that allowing superluminal speeds is not in direct contradiction with the special theory of relativity, which does not treat superluminality at all. The original derivation (Einstein 1905) of the theory of co-ordinate transformation is based on the definition of simultaneity enforcing the constancy of the speed of light. The synchronisation of clocks using light rays clearly cannot be done if the two frames are moving with respect to each other at superluminal speeds. All the ensuing equations apply only to subluminal speeds. It does not necessarily preclude the possibility of superluminal motion. However, an object starting from a fixed point and accelerating past the speed of light is clear violation of the special theory of relativity.

Another consequence of the traditional explanation of the apparent superluminal speed is that it is invariably associated with a blue shift. As given in equation (2), the apparent transverse superluminal speed, can occur only in a narrow region of $\cos \theta$. In this region, the longitudinal component of the velocity is always towards the observer, leading to a blue shift. The existence of blue shift associated with all superluminal jets has not been confirmed experimentally. Quasars with redshifts have been observed with associated superluminal jets. Two examples are: quasars 3C 279 (Wehrle et al. 2001) with a redshift $z = 0.536$ and 3C 216 with $z = 0.67$ (Paragi et al. 2000). However, the Doppler shift of spectral lines applies only to normal matter, not if the jets are made up of plasma, as currently believed. Thus, the current model of jets, made up of plasma collimated by a magnetic field originating from an accretion disc, can accommodate the lack of blue shift.

3. A Model for Symmetric Extragalactic Radio Sources

3.1. Symmetric Jets

Accepting hypothetical superluminal speeds, we can clearly tackle the second consequence of the traditional explanation (namely, the necessity of blue shift along with apparent su-

perluminal motion.) However, it is not clear how we perceive superluminal motion, because the light travel time effect will influence the way we perceive the kinematics. In this section, we will show that a single object moving superluminally, in a transverse direction across our field of vision, will look like two objects departing from a single point in a roughly symmetric fashion.




Fig. 4. The top panel (a) shows an object flying along $A'BA$ at a constant superluminal speed. The observer is at O . The object crosses B (the point of closest approach to O) at time $t = 0$. The bottom panel (b) shows how the object is perceived by the observer at O . It first appears at B' , then splits into two. The two apparent objects seem to go away from each other (along J_1 and J_2) as shown.

Consider an object moving at a superluminal speed as shown in Fig. 4(a). The point of closest approach is B . At that point, the object is a distance of y from the observer at O . Since the speed is superluminal, the light emitted by the object at some point B' (before the point of closest approach B) reaches the observer *before* the light emitted at A' . This gives an illusion of the object moving in the direction from B' to A' , while in reality it is moving in the opposite direction.

We use the variable t' to denote the the observer's time. Note that, by definition, the origin in the observer's time axis is set when the object appears at B . ϕ is the observed angle with respect to the point of closest approach B . ϕ is defined as $\theta - \pi/2$ where θ is the angle between the object's velocity and the observer's line of sight. ϕ is negative for negative time t .

It is easy to derive the relation between t' and ϕ . (See Appendix for the mathematical details.)

$$t' = y \left(\frac{\tan \phi}{\beta} + \frac{1}{\cos \phi} - 1 \right) \quad (3)$$

Here, we have chosen units such that $c = 1$, so that y is also the time light takes to traverse BO . The observer's time is measured with respect to y . i.e., $t' = 0$ when the light from the point of closest approach B reaches the observer.

The actual plot of ϕ as a function of the observer's time is given in Fig. 5. Note that for subluminal speeds, there is only one angular position for any given t' . The time axis scales with y . For subluminal objects, the observed angular position changes almost linearly with the observed time, while for superluminal objects, the change is parabolic.

Equation 3 can be approximated using a Taylor series expansion as:

$$t' \approx y \left(\frac{\phi}{\beta} + \frac{\phi^2}{2} \right) \quad (4)$$

From the quadratic equation 4, one can easily see that the minimum value of t' is $t'_{\min} = -y/2\beta^2$ and it occurs at $\phi_0 = -1/\beta$. Thus, to the observer, the object first appears (as though out of nowhere) at the position ϕ_0 at time $t'_{\min}$. Then it appears to stretch and split, rapidly at first, and slowing down later. This apparent time evolution of the object is shown in Fig. 9, where it is compared to the microquasar GRS 1915+105 (Mirabel & Rodríguez 1994; Fender et al. 1999).

The angular separation between the objects flying away from each other is:

$$\Phi = \frac{2}{\beta} \sqrt{1 + \frac{2\beta^2}{y} t'} = \frac{2}{\beta} (1 + \beta\phi) \quad (5)$$

And the rate at which the separation occurs is:

$$\frac{d\Phi}{dt'} = \sqrt{\frac{2}{y\Delta t'}} = \frac{2\beta}{y(1 + \beta\phi)} \quad (6)$$

where $\Delta t' = t' - t'_{\min}$, the apparent age of the symmetric object.

This discussion shows that a single object moving across our field of vision at superluminal speed creates an illusion of an object appearing at a certain point in time, stretching and splitting and then moving away from each other. This time evolution is given in equation 3, and illustrated in the bottom panel of Fig. 4(b). Note that the apparent time t' is reversed with respect to the real time t in the region A' to B' . An event that happens near B' appears to happen before an event near A' . Thus, the observer may see an apparent violation of causality, but it is just a part of the light travel time effect.

Fig. 6 shows the apparent width of a superluminal object as it evolves. The width decreases with time, along its direction of motion. (See the appendix for the mathematical details.) Thus, the appearance is that of two spherical objects appearing out of nowhere, moving away from each other, and slowly getting compressed into thinner and thinner ellipsoids and then almost disappearing.




Fig. 5. The apparent angular positions of an object travelling at different speeds at a distance y of one million light years from us. The angular positions (ϕ in radians) are plotted against the observer's time t' in years.




Fig. 6. The apparent angular positions and sizes of an object travelling at $\beta = 300$ at a distance y of one million light years from us. The angular positions (ϕ in arc seconds) are plotted against the observer's time t' in years. The shaded region represents the apparent angular spread of the object, with an assumed diameter of 500 light years.

If there are multiple objects, moving as a group, at roughly constant superluminal speed along the same direction, their appearance would be a series of objects appearing at the same angular position and moving away from each other sequentially, one after another. The apparent knot in one of the jets always has a corresponding knot in the other jet.

The calculation presented in this article is done in two dimensions. If we generalise to three dimensions, we can explain the precession observed in some systems. Imagine a cluster of objects, roughly in a planar configuration (like a spiral galaxy, for instance) moving together at superluminal speeds with respect to us. All these objects will have the points of closest approach to us in small angular region in our field of vision – this region is around the point of minimum distance between

the plane and our position. If the cluster is rotating (at a slow rate compared to the superluminal linear motion), then the appearance to us would be the apparent jets changing directions as a function of time. The exact nature of the apparent precession depends on the spatial configuration of the cluster and its angular speed.

If we can measure the angle ϕ_0 between the apparent core and the point of closest approach, we can directly estimate the real speed of the object β . We can clearly see the angular position of the core. However, the point of closest approach is not so obvious. We will show in the next section that the point of closest approach corresponds to zero red shift. (This is obvious intuitively, because at the the point of closest approach, the longitudinal component of the velocity is zero.) If this point (ϕ_0) can be estimated accurately, then we can measure the speed directly, from the relation $\phi_0 = -1/\beta$.

3.2. Red shifts of the Hot Spots

We can also work out time evolution of the red shift of the hot spots. However, as relativistic Doppler shift equation is not defined for superluminal speeds, we need to work out the relationship between the redshift (z) and the speed (β) from first principles. This is easily done (see the appendix for the mathematical details):

$$1 + z = |1 - \beta \cos \theta| = |1 + \beta \sin \phi| \quad (7)$$

There are two solutions for z , corresponding to the apparent objects at the two different positions. However, as shown in the appendix, they are nearly identical for any given value of the observer's time, t' .

Since we allow superluminal speeds in our model of extragalactic radio sources, we can explain the radio frequency spectra of the hot spots as extremely red-shifted blackbody radiation. The β s involved in this explanation are typically very large, and we can approximate the red shift as:

$$1 + z \approx |\beta \phi| \approx \frac{|\beta \Phi|}{2} \quad (8)$$

Assuming the object to be a black body similar to the sun, we can predict the peak wavelength (defined as the wavelength at which the luminosity is a maximum) of the hot spots as:

$$\lambda_{\max} \approx (1 + z)480\text{nm} \approx \frac{|\beta \Phi|}{2}480\text{nm} \quad (9)$$

where Φ is the angular separation between the two hot spots.

This shows that the peak RF wavelength increases linearly with the angular separation. If multiple hot spots can be located in a twin jet system, their peak wavelengths will depend only on their angular separation, in a linear fashion. The real speed of the single object masquerading as two hot spots can be estimated from peak wavelength measurements. Furthermore, if the measurement is done at a single radio frequency, intensity variation can be expected as the hot spot moves along the jet.

Fig. 7 shows the variation of red shift as a function of the observer's time. In the top panel, we have plotted an object with $\beta = 300$ and $y =$ one million light years. For $t' < 0$, there is a strong blue shift, which explains the observed, transient




Fig. 7. Time evolution of the red shift from a superluminal object. The top panel (a) shows the red shifts expected from an object moving at $\beta = 300$ at a distance of one million light years from us. In the bottom panel (b), the object is moving at a speed $\beta = 300\,000$ at a distance of 15 billion light years. Note that the two apparent objects have nearly identical red shifts.

hard X-ray spectra of some of the symmetric jets. Donley et al. (2002) have a recent survey of such data, though the currently favoured explanation for such transient emissions is a stellar tidal disruption scenario. The small difference between the red shifts of the two apparent objects may explain the double peak structure observed in the spectral data of some of the AGNs (Eracleous & Halpern 2003).

In order to have a red shift that will push a black body radiation of a star similar to our sun into RF regions would require a β of about 300 000 and a distance of closest approach of about 15 billion light years. This is plotted in the bottom panel of Fig. 7.

3.3. Summary of Predictions

Some of the different quantitative predictions of the model are recapitulated here. These are predictions that are relatively easy to verify with existing data.

- The appearance of a single object moving across our field of vision at superluminal speed is that of an object appearing at a point, splitting and moving away in opposite directions.
- The core will always have a fixed angular position.

- The new superluminal knots appearing in the jets will always appear in pairs.
- The two apparent objects will shrink monotonically. As the knots move away from the core, they become thinner and thinner ellipsoids, contracting along the direction of motion while the trasverse size remains roughly constant.
- The separation speed is very high in the beginning, but it slows down parabolically with time.
- The hot spots have almost identical RF spectra (and red shifts).
- The RF wavelength at which the luminosity of the hot spots is the maximum increases linearly with the angular separation between them.
- Close to the core, the the spectrum is heavily blue shifted. Thus, the object can be a strong X -ray or even γ ray source for a brief period of time. After that, the spectrum moves through optical to RF region.

A clear indication of a movement in the core's angular position, or a superluminal knot appearing without a counterpart in the opposite jet will be strong evidence against our model based on superluminality. On the other hand, a clear measurement of apparent superluminal motion in both the jets (not reported so far) will provide a convincing indication that the conventional explanation is inadequate.

3.4. Comparison to Measurements




Fig. 8. The angular speed of M87 (as a function of the feature angle ϕ) fitted against our model. The solid line constrains the distance of closest approach to the estimated distance of M87 – 52 million light years, giving a $\beta = 4.8$. The dotted line is a free fit giving $\beta = 84\,000$.

Biretta et al. (1999) have reported proper motion in one of the jets of M87 as a function of the angle (ϕ) between the apparent core and the feature points. M87 is estimated to be at about 52 million light years away from us, which gives us the value of y . In equation 6, we have the apparent angular speed ($d\Phi/dt'$) as a function of ϕ . Making the reasonable approximation $\Phi \approx 2\phi$, we can fit these data to our equation. The result is shown in Fig. 8. The fit gives a value $\beta = 4.8$. A cluster of objects flying across our field of vision, about five times

faster than light and at a distance of 52 million light years, will look like two jets moving away from each other at roughly 38 mas/year. If one of the two jets is hidden for some reason, the appearance will be a single jet of objects moving away from a point with an angular speed of about 19 mas/year. Note that we exclude the first two points from the fit. In this region close to the core, the appearance of new objects makes it difficult to track the features.




Fig. 9. Fit of our model to the time evolution measurements of GRS1915+105. The yellow curve overlaid corresponds to our perception of a single superluminal object, travelling at $\beta = 3\,000$ across our field of vision.

A much better fit can be obtained if we were to let the distance y also float. The resulting β of about 90 000 may explain the spectra of the hot spots. While the estimated β may look excessive, once superluminal motion is allowed, there is no a priori reason why it should not take any value at all. Fig. 9 shows another comparison of our model to the data available in the literature. Here, the time evolution of the microquasar GRS 1915+105 (Mirabel & Rodríguez 1994; Fender et al. 1999) is fitted to our model of a single superluminal object. The deceleration of relativistic jets (one of the predictions from our model) has been observed in the Microquasar XTE J1550-564 (Corbel et al. 2002), though it is currently believed to be an effect similar to frictional drag.

AGNs are known to have intensely blue or ultraviolet core, not easily explained by thermal models. But this is an expected feature in our model. As seen in equation 9, the core (where $\Phi \rightarrow 0$) must have a highly blue shifted spectrum. A clear evolution of emission frequency from ultraviolet to RF is seen in the photometry of the jet in 3C 273 (Jester et al. 2004). The spectrum shifts from lower wavelength to higher as a function of the angular distance from the core, strikingly consistent with our prediction.

This shifting of peak frequency can be seen at a much larger scale in Fig. 2. The size of the optical core is about a tenth of the angular separation between the hotspots. If we model Cygnus A as a collection of objects moving together at superluminal speeds, the core region would have emissions in the γ , X-ray, UV or optical region. As we move away towards the hotspots, the peak frequency would continuously shift to RF. This behaviour is indeed reported (Bach et al. 2004) recently. This also partially explains why extragalactic radio sources seem to be associated with galactic nuclei, instead of appearing randomly in the sky. A large collection of objects moving together (a large spiral galaxy, viewed from the side, for instance) superluminally gives the impression of a smaller stationary object with optical emission. The apparent object is likely to appear elongated along the direction of motion (with the major axis along the direction of the jets), with RF lobes appearing symmetrically farther away from the core. If the motion is not along a linear trajectory, we may see curved jets.

4. Conclusions

In this article, we explored the full implications of the traditional explanation for the apparent superluminal motion observed in certain quasars and microquasars. The equation that explains the apparent superluminal speeds predicts that objects receding from us should appear to be moving slower. Thus, in a symmetric radio sources where it is observed, the superluminal motion can appear only in one of the jets. The observed symmetry of these extragalactic radio sources (even subluminal ones) is incompatible with the explanation. Another consequence is that an apparent superluminal motion (if the moving objects are composed of normal matter rather than plasma) must always show a blue shift, a red shifted object can never be superluminal.

We further explored the possibility of real superluminal motion. We showed that a single superluminal object flying across our field of vision would appear to us as a symmetric separation of two objects from a fixed core. Using this fact as the model for such symmetric jets, we can explain their kinematical features quantitatively. In particular, we showed that the angle of separation of the hot spots is parabolic in time, and the red shifts of the two hot spots are almost identical to each other. Even the fact that the spectra of the hot spots are in the radio frequency region can be explained by assuming hyperluminal motion and the consequent red shift of the blackbody radiation. Furthermore, the requirement that an apparent superluminal motion be associated with a blue shift does not apply any more.

We presented a set of predictions and compared them to existing data. The features such as the blueness of the core, symmetry of the lobes, the transient γ and X-Ray bursts, the measured evolution of the spectra along the jet all find natural and simple explanations in this model. Note, however, that we have not addressed the dynamics of the model – how are the super or hyperluminal objects powered? The only observation in this article is that a collection of objects travelling superluminally across our field of vision can appear remarkably similar to an AGN with symmetrically placed radio frequency lobes and

hotspots. It does not preclude plasma jets that may be related to space-time singularities or other massive objects and the associated accretion discs. The conventional explanation of the apparent superluminal motion in asymmetric jets (e.g., quasar 3C 279 Wehrle et al. (2001)) also stands. In fact, our model is just a generalisation of the conventional explanation.

We argued that superluminal motion is not inconsistent with the special theory of relativity, which just does not deal with it. Acceptance of superluminality has far-reaching consequences in other long established notions of our universe. In particular, it can be shown (see Appendix B) that the apparent expansion of the universe at strictly subluminal speed is also an artifact of our perception of superluminal motion. Thus, the theory of the big bang will have to be looked at once again to see how light travel time effect modifies it. The description of extragalactic (or galactic) radio sources in terms of superluminal motion has a direct impact on our understanding of black holes.

Appendix A: Mathematical Details




Fig. A.1. The object is flying along AB , the observer is at O . The object crosses A at time $t = 0$. It reaches A' at time $t = t'$. A photon emitted at A reaches O at time $t = t_0$, and a photon emitted at A' reaches O at time $t = t''$.

A.1. Velocity Profile of an Expanding Object

This section re-derives the ellipse in Fig. 3 from first principles. In Fig. A.1, there is an observer at O . An object is flying by at a high speed $v = \beta c$ along the horizontal line AB . With no loss of generality, we can assume that $t = 0$ when the object is at A . It passes A' at time t' . The photon emitted at time $t = 0$ reaches the observer at time $t = t_0$, and the photon emitted at A' (at time $t = t'$) reaches him at time $t = t''$. The angle between the object's velocity at A and the observer's line of sight is θ . We have the Pythagoras equations:

$$z^2 = x^2 + y^2 \quad \& \quad z'^2 = x'^2 + y^2 \quad (\text{A.1})$$

$$\Rightarrow \frac{x + x'}{z + z'} = \frac{z - z'}{x - x'} \quad (\text{A.2})$$

If we assume that x and z (distances at time t_0) are not very different from x' and z' respectively (distances at time t'), we can write,

$$\cos \theta = \frac{x}{z} \approx \frac{x + x'}{z + z'} = \frac{z - z'}{x - x'} \quad (\text{A.3})$$

We define the real speed of the object as:

$$v = \beta c = \frac{x - x'}{t'} \quad (\text{A.4})$$

But the speed it *appears* to have will depend on when the observer senses the object at A and A'. The apparent speed of the object is:

$$v' = \beta' c = \frac{x - x'}{t'' - t_0} \quad (\text{A.5})$$

Thus,

$$\frac{\beta}{\beta'} = \frac{t'' - t_0}{t'} \quad (\text{A.6})$$

$$= 1 + \frac{z' - z}{ct'} \quad (\text{A.7})$$

$$= 1 - \frac{x - x'}{ct'} \cos \theta \quad (\text{A.8})$$

$$= 1 - \beta \cos \theta \quad (\text{A.9})$$

which gives,

$$\beta' = \frac{\beta}{1 - \beta \cos \theta} \quad (\text{A.10})$$

Fig. 3 is the locus of β' for a constant $\beta = 0.8$, plotted against the angle θ .

A.2. Superluminal Red shift

Red shift (z) defined as:

$$1 + z = \frac{\lambda'}{\lambda} \quad (\text{A.11})$$

where λ' is the measured wavelength and λ is the known wavelength. In Fig. A.1, the number of wave cycles created in time t' between A and A' is the same as the number of wave cycles sensed at O between t_0 and t'' . Substituting the values, we get:

$$\frac{t' c}{\lambda} = \frac{(t'' - t_0) c}{\lambda'} \quad (\text{A.12})$$

Using the definitions of the real and apparent speeds, it is easy to get

$$\frac{\lambda'}{\lambda} = \frac{\beta}{\beta'} \quad (\text{A.13})$$

Using the relationship between the real speed β and the apparent speed β' (equation (A.10)), we get

$$1 + z = \frac{1}{1 + \beta' \cos \theta} = 1 - \beta \cos \theta \quad (\text{A.14})$$

As expected, z depends on the longitudinal component of the velocity of the object. Since we allow superluminal speeds in this calculation, we need to generalise this equation for z noting that the ratio of wavelengths is positive. Taking this into account, we get:

$$1 + z = \left| \frac{1}{1 + \beta' \cos \theta} \right| = |1 - \beta \cos \theta| \quad (\text{A.15})$$




Fig. A.2. An object flying along A'BA at a constant superluminal speed. The observer is at O. The object crosses B (the point of closest approach to O) at time $t = 0$.

A.3. Kinematics of Superluminal Objects

The derivation of the kinematics is based on Fig. A.2. Here, an object is moving at a superluminal speed along A'BA. At the point of closest approach, B, the object is a distance of y from the observer at O. Since the speed is superluminal, the light emitted by the object at some point B' (before the point of closest approach B) reaches the observer *before* the light emitted at A'. This gives an illusion of the object moving in the direction from B' to A', while in reality it is moving from A' to B'. ϕ is the observed angle with respect to the point of closest approach B. ϕ is defined as $\theta - \pi/2$ where θ is the angle between the object's velocity and the observer's line of sight. ϕ is negative for negative time t .

We choose units such that $c = 1$, in order to make algebra simpler. t' denotes the observer's time. Note that, by definition, the origin in the observer's time, t' is set when the object appears at B.

The real position of the object at any time t is:

$$x = y \tan \phi = \beta t \quad (\text{A.16})$$

A photon emitted at t will reach O after traversing the hypotenuse. A photon emitted at B will reach the observer at $t = y$, since we have chosen $c = 1$. If we define the observer's time t' such that the time of arrival is $t = t' + y$, then we have:

$$t' = t + \frac{y}{\cos \phi} - y \quad (\text{A.17})$$

which gives the relation between t' and ϕ .

$$t' = y \left(\frac{\tan \phi}{\beta} + \frac{1}{\cos \phi} - 1 \right) \quad (\text{A.18})$$

Expanding the equation for t' to second order, we get:

$$t' = y \left(\frac{\phi}{\beta} + \frac{\phi^2}{2} \right) \quad (\text{A.19})$$

The minimum value of t' occurs at $\phi_0 = -1/\beta$ and it is $t'_{\min} = -y/2\beta^2$. To the observer, the object first appears at the position $\phi = -1/\beta$. Then it appears to stretch and split, rapidly at first, and slowing down later. The angular separation between the

objects flying away from each other is the difference between the roots of the quadratic equation (A.19):

$$\Phi = \phi_1 - \phi_2 = \frac{2}{\beta} \sqrt{1 + \frac{2\beta^2}{y} t'} = \frac{2}{\beta} (1 + \beta\phi) \quad (\text{A.20})$$

We also have the mean of the roots equal to the position of the minimum:

$$\phi_1 + \phi_2 = \frac{-2}{\beta} \quad (\text{A.21})$$

And the rate at which the separation occurs is:

$$\frac{d\Phi}{dt'} = \frac{2\beta}{y \sqrt{1 + \frac{2\beta^2}{y} t'}} = \frac{2\beta}{y(1 + \beta\phi)} \quad (\text{A.22})$$

Defining $t'_0 = -t_{\min}$, we can write:

$$\frac{d\Phi}{dt'} = \sqrt{\frac{2}{y\Delta t'}} \quad (\text{A.23})$$

where $\Delta t' = t' + t'_0$, the apparent age of the quasar.

As shown before in equation (A.15), the red shift z depends on the real speed β as:

$$1 + z = |1 - \beta \cos \theta| = |1 + \beta \sin \phi| \quad (\text{A.24})$$

There are two solutions for z . For $\sin \phi < -1/\beta$, you get

$$z_1 = -2 - \beta \sin \phi_1 \quad (\text{A.25})$$

and for $\sin \phi > -1/\beta$,

$$z_2 = \beta \sin \phi_2 \quad (\text{A.26})$$

Thus, we get the difference in the red shift between the two hot spots as:

$$\Delta z \approx 2 + \beta(\phi_1 + \phi_2) \quad (\text{A.27})$$

But $\phi_1 + \phi_2 = -2/\beta$ and hence $\Delta z = 0$. The two hot spots will have identical red shifts, if terms of ϕ^3 and above are ignored.

A.4. Time Evolution of Object Size and Red Shift

Fig. 6 shows the apparent positions (ϕ) and the size of the superluminal object as the observer sees it, as a function of the observer's time (t'). Fig. 7 is a similar time evolution of the red shift (z). In this section, we describe how these two plots are created. It is easiest to express the quantities parametrically as a function of the real time t . Referring to Fig. A.2, we write,

$$x = \beta t \quad (\text{A.28})$$

$$t' = t + \sqrt{\beta^2 t^2 + y^2} - y \quad (\text{A.29})$$

$$\sin \phi = \frac{\beta t}{\sqrt{\beta^2 t^2 + y^2}} \quad (\text{A.30})$$

The solid parabola in Fig. 6 is ϕ vs. t' from these equations as t is varied between -40 and 20 years, with $y = 1\,000\,000$ light years and $\beta = 300$.

In order to get the variation of the size of the object (the shaded region in Fig. 6), we assume a diameter $d = 500$ light years.

$$t'_\pm = t + \sqrt{\left(\beta t \pm \frac{d}{2}\right)^2 + y^2} - y \quad (\text{A.31})$$

$$\sin \phi_\pm = \frac{\beta t \pm \frac{d}{2}}{\sqrt{\left(\beta t \pm \frac{d}{2}\right)^2 + y^2}} \quad (\text{A.32})$$

The boundaries of the shaded region are given by ϕ_+ vs. t'_+ and ϕ_- vs. t'_- .

As shown before (see equation (A.24), the red shift z depends on the real speed β as:

$$1 + z = |1 + \beta \sin \phi| = \left| 1 + \frac{\beta^2 t}{\sqrt{\beta^2 t^2 + y^2}} \right| \quad (\text{A.33})$$

Since we know z and t' functions of t , we can plot their interdependence parametrically. This is shown in Fig. 7, for two sets of values for β and y .

Appendix B: Apparent Expansion of the Universe

The apparent recessional speed is the longitudinal component of β' is $\beta'_\parallel = \beta' \cos \theta$. From equation (1), we can see that

$$\lim_{\beta \rightarrow \pm\infty} \beta'_\parallel = -1 \quad (\text{B.1})$$

The apparent recessional speed (which can be measured using red shifts) tends to c (or, $\beta'_\parallel \rightarrow -1$), when the real speed is highly superluminal. This limit is independent of the actual direction of motion of the object θ . Thus, whether a superluminal object is receding or approaching (or, in fact, moving in any other direction), the appearance from our perspective would be an object receding roughly at the speed of light. This appearance of all (possibly superluminal) objects receding from us at strictly subluminal speeds is an artifact of our perception, rather than the true nature of the universe.

References

- Bach, U., Krichbaum, T. P., Middelberg, E., et al. 2004, in Proceedings of the 7th European VLBI Network Symposium, ed. R. Bachiller, F. Colomer, & P. Desmurs, J.F. and de Vicente
- Biretta, J. A., Sparks, W. B., & Macchetto, F. 1999, ApJ, 520, 621
- Corbel, S., Fender, R. P., Tzioumis, A. K., et al. 2002, Science, 298, 196
- Donley, J. L., Brandt, W. N., Eracleous, M. C., & Boller, T. 2002, AJ, 124, 1308
- Einstein, A. 1905, Annalen der Physik, 17, 891
- Eracleous, M. & Halpern, J. 2003, ApJ, 599, 886
- Fender, R. P., Garrington, S. T., McKay, D. J., et al. 1999, MNRAS, 304, 865
- Gisler, G. 1994, Nature, 371, 18
- Jester, S., Roeser, H. J., Meisenheimer, K., & Perley, R. 2004, A&A (in press)

- Laing, R. A., Parma, P., de Ruiter, H. R., & Fanti, R. 1999, MNRAS, 306, 513
- Mirabel, I. F. & Rodríguez, L. F. 1994, Nature, 371, 46
- Mirabel, I. F. & Rodríguez, L. F. 1999, ARA&A, 37, 409
- Owsianik, I. & Conway, J. E. 1998, A&A, 337, 69
- Paragi, Z., Frey, S., Fejes, I., et al. 2000, PASJ, 52, 983
- Perley, R. A., Dreher, J. W., & Cowan, J. J. 1984, ApJ, 285, L35
- Polatidis, A. G., Conway, J. E., & Owsianik, I. 2002, in Proceedings of the 6th European VLBI Network Symposium, ed. Ros, Porcas, Lobanov, Zensus
- Rees, M. 1966, Nature, 211, 468
- Wehrle, A. E., Piner, B. G., Unwin, S. C., et al. 2001, ApJS, 133, 297
- Zensus, A. J. 1997, ARA&A, 35, 607